



Guidance for Substantiating the Evidence for Beneficial Effects of Probiotics: Prevention and Management of Infections by Probiotics^{1–3}

Danielle Wolvers,⁴ Jean-Michel Antoine,⁵ Eveliina Myllyluoma,⁶ Juergen Schrezenmeir,⁷ Hania Szajewska,⁸ and Ger T. Rijkers⁹

⁴Unilever Food and Health Research Institute, Vlaardingen, The Netherlands; ⁵Danone Research Center, Palaiseau, France; ⁶Institute of Biomedicine, Pharmacology, University of Helsinki and Valio Research Centre, Valio Ltd., Helsinki, Finland; ⁷Max Rubner-Institut, Karlsruhe, Germany; ⁸Department of Paediatrics, The Medical University of Warsaw, Warsaw, Poland; and ⁹Department of Surgery, University Medical Center Utrecht, and Department of Medical Microbiology and Immunology, St. Antonius Hospital, Nieuwegein, The Netherlands

Abstract

The rationale for the use of probiotics in the management of infectious diseases is supported by their potential to influence and stabilize the composition of gut microbiota, enhance colonization resistance, and modulate immune function parameters. A literature review was conducted to determine the efficacy of using probiotics in selected infections: 1) infectious diarrhea in infants and children, 2) traveler's diarrhea, 3) necrotizing enterocolitis in infants, 4) *Helicobacter pylori* infection, 5) respiratory tract infections in adults and children, 6) ear, nose, and throat infections, and 7) infectious complications in surgical and critically ill patients. The different types of infections that have been subject to clinical studies with different probiotics obviously prevent any generic conclusions. Furthermore, the lack of consistency among studies focusing on 1 specific infection, in study design, applied probiotic strains, outcome parameters, and study population, along with the still limited number of studies, preclude clear and definite conclusions on the efficacy of probiotics and illustrate the need for better-aligned study designs and methodology. Exceptions were the management of infectious diarrhea in infants and traveler's diarrhea, antibiotic-associated diarrhea, and necrotizing enterocolitis. Sufficient consistent data exist for these applications to conclude that certain probiotics, under certain conditions, and in certain target populations, are beneficial in reducing the risk of infection. In addition, some evidence exists, although conclusions are premature, for the management of *Helicobacter pylori* infection and possible reduction of treatment side effects. Certain probiotics may also reduce the risk of various symptoms of respiratory tract infections in adults and children, including ear, nose, and throat infections, although data are currently far too limited to distill any clinical recommendations in this area. Positive but also negative results have been obtained in prevention of infectious complications in surgical and critically ill patients. For future studies it is recommended that researchers provide adequate power, identify pathogens, and report both clinical outcomes and immune biomarkers relating to putative underlying mechanisms. *J. Nutr.* 140: 698S–712S, 2010.

Risk reduction and management of infections by probiotics

In 1916, Nissle demonstrated that transferring members of the human gut microbiota to healthy typhoid carriers resulted in

Salmonella being cleansed from their intestines (1). This finding pointed for the first time to one of the most important

¹ Published in a supplement to *The Journal of Nutrition*. Presented at the workshop "Guidance for Assessing Probiotics Beneficial Effects: How to Fill the GAP," held in Montreux, Switzerland, May 22–24, 2008 and organized by ILSI Europe in association with the International Dairy Federation. The supplement coordinator for this supplement is Agnès Méheust, ILSI Europe. Publication costs for this supplement were defrayed in part by the payment of page charges. This publication must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact. Supplement Coordinator disclosure: Agnès Méheust is employed as a Scientific Project Manager for ILSI Europe, which is largely funded by the food and related industries. Supplement Guest Editor disclosure: Josef Neu declares no conflict of interest. The opinions expressed in this publication are those of the authors and are not attributable to the sponsors or the publisher, editor, or editorial board of *The Journal of Nutrition*.

² Author Disclosures: This work was commissioned by the Probiotics Task Force of ILSI Europe composed of the following industry members: Barilla G. & R. Fratelli, Danisco, Danone, Friesland Campina, Kraft Foods, Mead Johnson Nutrition, Nestlé, Seven Seas, Unilever, Valio and Yakult Europe. H. Szajewska and G.T. Rijkers have no conflicts of interest to declare. J.-M. Antoine is an employee of Danone. E. Myllyluoma is an employee of Valio. J. Schrezenmeir received grants and/or consulting agreement and/or honorarium as speaker from Bauer, Boehringer Ingelheim, Campina, Danisco, Danone, Chr. Hansen, Infectopharm, Merck, Mona, Morinaga, Müller, Nestlé, NÖM, Orthomol, Pharmatech, Wakunaga, and Yakult. D.W. is an employee of Unilever. The opinions expressed herein are those of the authors and do not necessarily represent the views of ILSI Europe.

³ Supplemental Tables 1–13 are available as Online Supporting Material with the online posting of this paper at jn.nutrition.org. Correspondence should be addressed to ILSI Europe. E-mail: publications@ilsieurope.be.

physiological functions of the gut microbiota, i.e., to defend the gut against the colonization by exogenous microorganisms, now termed colonization resistance. Since that time, it has been demonstrated that different aerobic as well as anaerobic species of the gut microbiota are involved in this function (2).

Our understanding of the role of the microbiota in the resistance against infections has further evolved, and the importance of the interaction of the microbiota with the immune system and intestinal epithelial cells providing additional barriers to infectious agents is increasingly being recognized (3).

The potential impact that probiotics may have at the level of gut microbiota, on gut epithelium and its associated mucosal immune system, as well as systemically (4) provides a rationale of why probiotics are promising food components for the reduction of risk or management of infectious diseases. Most notably these relate to infections of the gastrointestinal tract, but infections at other sites have also been targeted recently.

For our review of the existing literature we defined a number of categories of infections, based on the localization of the infection (respiratory tract, gastrointestinal, systemic) and target populations (children, adults, elderly; at-risk groups or patients). Studies dealing with the effect of probiotics on urinary tract, vaginal infections, and skin infections were not analyzed. Furthermore, although there are many indications that various immune function parameters may be affected by probiotics, we chose to focus on studies investigating a clinical outcome and did not incorporate studies that solely describe effects of probiotics on biomarkers of the immune system.

Based on the available information, the subcategories are the following:

1. Infectious diarrhea in infants and children including acute infectious diarrhea and antibiotic-associated diarrhea (AAD).¹⁰
2. Traveler's diarrhea (TD).
3. Necrotizing enterocolitis (NEC) in infants.
4. *Helicobacter pylori* infection.
5. Respiratory tract infections in adults and children.
6. Ear, nose, and throat (ENT) infections.
7. Infectious complications in surgical and critically ill patients.

Infectious diarrhea

Acute infectious diarrhea. Oral rehydration is the mainstay of therapy for acute gastroenteritis and should continue to be fostered, encouraged, and supported. However, despite its proven efficacy, oral rehydration therapy remains underused (5). The main reasons for this are that an oral rehydration solution does not reduce the frequency of bowel movements and fluid loss, nor does it shorten the duration of the illness, which reasons both decrease acceptance. As a result, there is an interest on the part of patients, caregivers, and practitioners in simple, safe, and effective measures that will visibly reduce the rate of stool loss and/or the duration of diarrhea.

The evidence from several meta-analyses of randomized controlled trials (RCT) (6–9) [Table 1 (6–74)] consistently shows significant effects and moderate clinical benefits of some probiotic strains in the management of acute watery diarrhea, often rotaviral, primarily in infants and young children. So far, the beneficial effects of probiotics in acute infectious diarrhea appear

to be moderate (~1-d reduction in the duration of diarrhea), strain-dependent (75), dose-dependent [greater for doses $>10^{10}$ – 10^{11} colony-forming units (CFU)/d], significant in watery diarrhea and viral gastroenteritis but absent in invasive, bacterial diarrhea, more evident when treatment with probiotics is initiated early in the course of disease, and more evident in developed countries.

The use of probiotics for acute infectious diarrhea in children is an accepted therapy in Europe. The European Society for Paediatric Gastroenterology, Hepatology and Nutrition and European Society of Paediatric Infectious Diseases Expert Working Group have stated that selected probiotics with proven clinical efficacy [e.g., *Lactobacillus rhamnosus* GG (76), *Saccharomyces boulardii* (77)] and in appropriate dosage, according to the strain and the population (Table 1) may be used as an adjunct for the management of children with acute gastroenteritis being given rehydration therapy (78). Other probiotic strains may also be used provided their efficacy is documented in high-quality RCTs (or in meta-analyses).

AAD and *Clostridium difficile* diarrhea

A common side effect of antibiotic treatment is AAD, defined as otherwise unexplained diarrhea occurring in association with antibiotics administration (79). In the pediatric population, AAD occurs in ~11–40% of children between the initiation of therapy and up to 2 mo after cessation of treatment (80,81).

Several systematic reviews with or without meta-analysis documented that most of the tested probiotics have been shown to be effective in reducing the risk of AAD in the general (mainly adult) population (28–31). Also, a few of the more recent trials in adults, not included yet in the meta-analyses, showed encouraging results in prevention of AAD (34,35). Evidence from 3 recent systematic reviews of RCTs also suggests that probiotics reduce the risk of AAD in children (25–27) (Table 1).

In contrast, there is only weak or inconclusive evidence from 2 systematic reviews for the effectiveness of probiotics for the prevention and treatment of *C. difficile*-associated diarrhea (Table 1). The authors of the first review concluded that available evidence (all in adults) does not support the administration of probiotics with antibiotics to prevent or treat *C. difficile* diarrhea (38). The conclusions from the second systematic review (with meta-analysis) do support probiotic use (82). However, the latter meta-analysis has been criticized for combining the results from 1 study on prevention of *C. difficile* diarrhea with results from 5 studies on treatment of *C. difficile* diarrhea and pooling data on different probiotics, different conditions, and different patient characteristics as well as some methodological issues calling for caution in interpreting the conclusions (82,83). In children, despite some anecdotal evidence of the efficacy of probiotics (84,85), no RCT investigating such a possibility has been conducted.

The current available evidence suggests a moderate beneficial effect of selected probiotics, such as *Saccharomyces boulardii* or *Lactobacillus rhamnosus* GG, in the prevention of AAD in children, or *Lactobacillus casei* DN 114 001 in the elderly. Because these probiotics have been shown to be capable of providing reasonable protection against the development of AAD, their use is probably warranted whenever preventing this usually self-limited complication is important. The available data provide evidence that selected probiotics significantly reduce the risk of diarrhea in patients treated with antibiotics in general. However, not all antibiotics are likely to be equal in causing AAD. Currently, conclusions about the efficacy of probiotics in preventing diarrhea attributable to any single antibiotic class cannot be made. Why some probiotics are

¹⁰ Abbreviations used: AAD, antibiotic-associated diarrhea; AOM, acute otitis media; AP, acute pancreatitis; CFU, colony-forming units; ENT, ear, nose, and throat; NEC, necrotizing enterocolitis; RCT, randomized controlled trial; TD, traveler's diarrhea.

TABLE 1 Probiotics in the prevention and treatment of infections¹

Disease/marker	Reference	Participants	Type of probiotic(s)	Duration	Type of study	Outcome
1. Respiratory tract						
Common cold	De Vrese et al. 2005 (10)	479 adults	<i>L. gasseri</i> , <i>B. longum</i> , <i>B. bifidum</i> mixture	3–5.5 mo	R, DB, PC	Reduction duration (1.9 d) No effect on incidence Reduction days with fever (0.7 d) Reduction symptoms
Common cold	Winkler et al. 2005 (11)	477 adults	<i>L. gasseri</i> , <i>B. longum</i> , <i>B. bifidum</i> mixture	3–5.5 mo	R, DB, PC	Reduction symptoms Reduction days with fever (54%) No effect on duration Near sign reduction in incidence (14%)
RT and GI symptoms	Tubelius et al. 2005 (12)	262 adults	<i>L. reuteri</i>	80 d	R, DB, PC	Reduced incidence sickness (>50%)
RT, ENT, and GI symptoms	Turchet et al. 2003 (13)	360 elderly	<i>L. casei</i>	3 wk	R, C	Reduced duration (1.7 d) No effect on incidence
RT and GI symptoms	Kekkonen et al. 2007 (14)	119 trained adults	<i>L. rhamnosus</i> GG	3 mo	R, DB, PC	No effect on incidence during training (intervention) Reduced duration (1.3 d) of GI symptoms during recovery (wash-out)
RT symptoms	Tiollier et al. 2007 (15)	47 trained adults	<i>L. casei</i>	1 mo	R, DB, PC	No effect on incidence, duration, and symptoms
RT symptoms	Cox et al. 2008 (16)	20 trained adults	<i>L. fermentum</i>	30 d	PC, DB, cross over	Reduction in number of days (>50%) Near sign reduction in symptoms
RT and GI symptoms	Hatakka et al. 2001 (17)	571 children	<i>L. rhamnosus</i> GG	7 mo (5 d/wk)	R, DB, PC	Reduced absence from daycare (0.9 d) Reduced incidence of complications (8.6%)
Respiratory illness, diarrhea, fever	Weizman et al. 2005 (18)	201 infants	<i>B. animalis</i> or <i>L. reuteri</i>	12 wk	R, DB, PC	Size of effects diminished after age adjustment
RT symptoms	Marsaglia et al. 2007 (19)	80 children	<i>B. clausii</i>	90 d	R, SB, C	Reduced incidence fever Reduced incidence and duration diarrhea No effect on RT
Otitis media	Hatakka et al. 2007 (20)	309 otitis-prone children and toddlers	Mixture of <i>L. rhamnosus</i> GG, <i>L. rhamnosus</i> LC705, <i>B. breve</i> 99, <i>P. freudenreichii</i> ssp. <i>shermanii</i> JS	6 mo	R, DB, PC	Reduced duration during treatment (2.7 d) and follow up (4.3 d)
Otitis media	Roos et al. 2001 (21)	108 otitis-prone children and toddlers	<i>Streptococcus mitis</i> , <i>Streptococcus sanguis</i> , <i>Streptococcus oralis</i> mixture (nasal spray)	10 d	R, DB, PC	Increased presence <i>M. Catarrhalis</i>
Tonsillitis	Roos et al. 1993 (22)	36 adults with recurrent tonsillitis	<i>S. sanguis</i> and <i>S. mitis</i> mixture (oral spray)	10 d	R, DB, PC	Reduced incidence of recurrence (22% vs. 42%)
Tonsillitis	Roos et al. 1996 (23)	112 adults with recurrent tonsillitis	<i>S. sanguis</i> and <i>S. mitis</i> mixture (oral spray)	10 d	R, DB, PC	Reduced incidence of recurrence (2% vs. 23%)
Tonsillitis	Falck et al. 1999 (24)	342 adults and children with recurrent tonsillitis	<i>S. sanguis</i> and <i>S. mitis</i> mixture (oral spray)	10 d	R, DB, PC	No effect on early recurrence Reduction of late recurrence
2. Probiotics in treatment of acute infectious diarrhea						
Diarrhea lasting 3 d	Szajewska et al. 2001 (6)	731 children	<i>L. rhamnosus</i> GG; <i>L. reuteri</i> ; <i>L. acidophilus</i> LB; <i>Streptococcus thermophilus</i> lactis; <i>L. acidophilus</i> and <i>L. bulgaricus</i> ; <i>S. boulardii</i> .	NA	R, PC	Relative risk (95% CI): 0.4 (0.3 to 0.5) Number needed to treat (95% CI): 4 (3 to 9)

(Continued)

TABLE 1 Continued

Disease/marker	Reference	Participants	Type of probiotic(s)	Duration	Type of study	Outcome
Diarrhea lasting 3 d	Allen et al. 2004 (9)	1341 children and adults	<i>L. rhamnosus</i> GG, <i>L. reuteri</i> , <i>L. acidophilus</i> LB, <i>Strep. thermophilus</i> lactis, <i>L. acidophilus</i> , and <i>L. bulgaricus</i> , <i>Enterococcus</i> SF68, <i>L. acidophilus</i> , and <i>L. bifidus</i> , <i>L. casei</i> , <i>L. rhamnosus</i> , and <i>L. reuteri</i> , <i>S. boulardii</i> .	NA	R, PC	Relative risk (95% CI): 0.7 (0.6 to 0.8) Number needed to treat (95% CI): 5 (4 to 7)
Duration of diarrhea	Szajewska et al. 2001 (6)	731 children	<i>L. rhamnosus</i> GG, <i>L. reuteri</i> , <i>L. acidophilus</i> LB, <i>Strep. thermophilus</i> lactis, <i>L. acidophilus</i> and <i>L. bulgaricus</i> , <i>S. boulardii</i> .	NA	R, PC	WMD (95% CI): -18 h (-27 to -10)
Duration of diarrhea	Van Niel et al. 2002 (7)	675 children	<i>L. rhamnosus</i> GG, <i>L. reuteri</i> , <i>L. acidophilus</i> , and <i>L. bulgaricus</i> .	NA	R, PC	WMD (95% CI): -17 h (-29 to -7)
Duration of diarrhea	Huang et al. 2002 (8)	1917 children	<i>L. rhamnosus</i> GG, <i>L. acidophilus</i> , <i>L. bulgaricus</i> , <i>S. thermophilus</i> , <i>L. rhamnosus</i> , <i>Yalacta</i> (<i>L. rhamnosus</i> , <i>L. delbrueckii</i> , <i>L. bulgaricus</i>), <i>L. reuteri</i> , <i>Enterococcus</i> SF68, <i>S. boulardii</i> , <i>S. subtilis</i> , <i>B. bifidum</i> and <i>B. infantis</i> .	NA	R, DB, PC, open label	WMD (95% CI): -19 h (-26 to -14)
Duration of diarrhea	Allen et al. 2004 (9)	970 children and adults	<i>L. rhamnosus</i> GG, <i>L. reuteri</i> , <i>L. acidophilus</i> LB, <i>Streptococcus thermophilus</i> lactis, <i>L. acidophilus</i> and <i>L. bulgaricus</i> , <i>Enterococcus</i> SF68, <i>L. acidophilus</i> and <i>L. bifidus</i> , <i>L. casei</i> , <i>L. rhamnosus</i> , and <i>L. reuteri</i> , <i>S. boulardii</i> .	NA	R, PC	WMD (95% CI): -30 h (-42 to -19)
3. Probiotics in prevention of AAD						
Prevention of AAD	Szajewska et al. 2006 (25)	766 children	<i>L. acidophilus</i> and <i>L. bulgaricus</i> (1 RCT); <i>L. acidophilus</i> and <i>B. infantis</i> (1 RCT); <i>B. lactis</i> Bb12 and <i>Str. thermophilus</i> (1 RCT); <i>L. rhamnosus</i> GG (2 RCTs); <i>S. boulardii</i> (1 RCT).	NA	R, PC	Relative risk (95% CI): 0.44 (0.25 to 0.77)
Prevention of AAD	Johnston et al. 2006 (26)	707 children	<i>L. acidophilus</i> and <i>L. bulgaricus</i> (1 RCT); <i>L. acidophilus</i> and <i>B. infantis</i> (1 RCT); <i>L. rhamnosus</i> GG (2 RCTs); <i>S. boulardii</i> (1 RCT); <i>L. reuteri</i> , <i>S. boulardii</i> , <i>S. sporogens</i> and fructooligosaccharides (1 RCT).	NA	R, PC	Relative risk (95% CI): 0.43 (0.25 to 0.75) Relative risk (95% CI): 1.01 (0.64 to 1.61)
Prevention of AAD	Johnston et al. 2007 (27)	1946 children	<i>L. rhamnosus</i> GG (2 RCTs); <i>S. boulardii</i> (3 RCTs); <i>B. lactis</i> and <i>Str. thermophilus</i> (1 RCT); <i>L. acidophilus</i> and <i>B. infantis</i> (1 RCT); <i>L. sporogenes</i> and fructooligosaccharides (1 RCT); <i>L. acidophilus</i> and <i>L. bulgaricus</i> (1 RCT).	NA	R, PC	Relative risk (95% CI): 0.49 (0.32 to 0.74)

(Continued)

TABLE 1 Continued

Disease/marker	Reference	Participants	Type of probiotic(s)	Duration	Type of study	Outcome
Prevention of AAD	D'Souza et al. 2002 (28)	830 children and adults	<i>S. boulardii</i>	NA	R, PC	OR (95% CI): 0.39 (0.25 to 0.62) Number needed to treat (95% CI): 11 (8 to 20)
		384 children and adults	<i>L. acidophilus</i> and <i>L. bulgaricus</i> (2 RCTs); <i>L. acidophilus</i> and <i>B. longum</i> (1 RCT); <i>L. rhamnosus</i> GG (1 RCT); <i>E. faecium</i> SF68 (1 RCT).			OR (95% CI): 0.34 (0.19 to 0.61) Number needed to treat (95% CI): 11 (8 to 18)
Prevention of AAD	Cremonini et al. 2002 (29)	Total: 1214 446 children and adults	Total <i>L. rhamnosus</i> GG (3 RCTs), <i>Lactobacillus</i> spp. (1 RCT)	NA	R, PC	OR (95% CI): 0.37 (0.26 to 0.53) Relative risk (95% CI): 0.5 (0.4 to 0.7) Number needed to treat (95% CI): 9 (7 to 17)
		435 children and adults	<i>S. boulardii</i>			Relative risk (95% CI): 0.6 (0.4 to 0.9) Number needed to treat (95% CI): 14 (8 to 18)
Prevention of AAD	Szajewska et al. 2005 (30)	Total: 881 1076 children and adults	<i>S. boulardii</i>	NA	R, PC	Relative risk (95% CI): 0.4 (0.3 to 0.6) Number needed to treat (95% CI): 9 (7 to 14)
Prevention of AAD	Hawrelak et al. 2005 (31)	692 children and adults	<i>L. rhamnosus</i> GG	NA	R, PC	No statistical pooling
Prevention of AAD	Sazawal et al. 2006 (32)	No data	Various	NA	R, PC	Relative risk (95% CI): 0.52 (0.35 to 0.65)
Prevention of AAD RCT	McFarland et al. 2006 (33)	2810 N (Exp/Cont)	Various	NA	R, PC	Relative risk (95% CI): 0.43 (0.31 to 0.58)
Prevention of AAD	Wenus et al. 2008 (34)	87 children and adults (46/41) FU 63/87 (72%)	<i>L. rhamnosus</i> GG, La-5, Bb-12	14 d	R, PC	Relative risk (95% CI): 0.21 (0.05 to 0.93)
Prevention of AAD	Hickson et al. 2007 (35)	113 children and adults (57/56) FU 113/135 (83%)	<i>L. casei</i> DN-114001, <i>S. thermophilus</i> , <i>L. bulgaricus</i>	Ab + 7 d	R, PC	Relative risk (95% CI): 0.36 (0.2 to 0.76) Number needed to treat (95% CI): 5 (3 to 16)
Prevention of AAD	Ruszczynski et al. 2008 (36)	240 children and adults (120/120)	<i>L. rhamnosus</i> (strains E/N, Oxy and Pen)	Ab	R, PC	Relative risk (95% CI): 0.45 (0.2 to 0.95) Number needed to treat (95% CI): 11 (6 to 106)
Prevention of AAD	Szymanski et al. 2008 (37)	78 children and adults (40/30)	<i>B. longum</i> PLO3, <i>L. rhamnosus</i> KL53A, <i>L. plantarum</i> PLO2	Ab	R, PC	Relative risk (95% CI): 0.47 (0.04 to 5) Number needed to treat (95% CI): Not significant
4. Probiotics in prevention and management of <i>C. difficile</i> diarrhea						
Treatment of <i>C. difficile</i> diarrhea	Dendukuri et al. 2005 (38)	No data	<i>S. boulardii</i> (2 RCTs), <i>L. plantarum</i> 299v (1 RCT)	NA	R, PC	No statistical pooling
Treatment of <i>C. difficile</i> diarrhea	McFarland et al. 2006 (33)	354 adults	<i>S. boulardii</i> (2 RCTs), <i>Lactobacillus</i> GG (2 RCTs), <i>L. plantarum</i> 299v (1 RCT), <i>L. acidophilus</i> and <i>B. bifidum</i> (1 RCT).	NA	R, PC	Relative risk (95% CI): 0.59 (0.4 to 0.85) Number needed to treat (95% CI): 8 (6 to 22)
Prevention of <i>C. difficile</i> diarrhea	Dendukuri et al. 2005 (38)	No data	<i>S. boulardii</i> (3 RCT), <i>Lactobacillus</i> GG (1 RCT), <i>L. acidophilus</i> and <i>B. bifidum</i> (1 RCT), <i>L. acidophilus</i> and <i>B. bifidum</i> (1 RCT), <i>L. acidophilus</i> and <i>B. bifidum</i>	NA	R, PC	No statistical pooling
Prevention of <i>C. difficile</i> diarrhea	McFarland et al. 2006 (33)	No data	<i>L. acidophilus</i> and <i>B. bifidum</i>	NA	R, PC	Relative risk (95% CI): 0.33 (0.07 to 1.59) Number needed to treat (95% CI): Not significant
5. Probiotics for prevention of NEC						
Prevention of NEC	Deshpande et al. 2007 (39)	1393 neonates	<i>B. breve</i> , <i>L. rhamnosus</i> GG; <i>S. boulardii</i> ; <i>B. infantis</i> and <i>S. thermophilus</i> and <i>B. bifidum</i> ; <i>L. acidophilus</i> and <i>B. infantis</i> ; <i>Lactobacillus casei</i> ; <i>B. lactis</i> .	3 - > 6 wk	R, PC	Outcome measure: NEC stage 2 Relative risk (95% CI): 0.36 (0.2 to 0.65) Number needed to treat (95% CI): 30 (19 to 71)

(Continued)

TABLE 1 Continued

Disease/marker	Reference	Participants	Type of probiotic(s)	Duration	Type of study	Outcome
Prevention of NEC	AlFaleh et al. 2008 (40)	1264 neonates	<i>B. infantis</i> and <i>S. thermophilus</i> and <i>B. bifidum</i> ; <i>S. boulardii</i> ; <i>L. rhamnosus</i> GG; <i>L. acidophilus</i> and <i>B. infantis</i> ; <i>L. casei</i> .	14 d- discharge	R, PC	Outcome measure: NEC stage 2 Relative risk (95% CI): 0.32 (0.17 to 0.60) Number needed to treat (95% CI): 26 (17 to 57)
Prevention of necrotizing enterocolitis	Samanta et al., 2009 (41)	91+95 neonates	<i>B. infantis</i> and <i>B. bifidum</i> and <i>B. longum</i> and <i>L. acidophilus</i> .	Until discharge	R, PC	Outcome measure: NEC stage 2 Relative risk (95% CI): 0.48 (0.22 to 1.04) Number needed to treat (95% CI): 10 (5 to 63)
Prevention of NEC	Lin et al. 2008 (42)	217+217 neonates	<i>B. bifidum</i> and <i>L. acidophilus</i>	14 d- discharge	R, PC	Outcome measure: NEC stage 2 Relative risk (95% CI): 0.43 (0.19 to 1.00) Number needed to treat (95% CI): 22 (12 to 113)
6. Probiotics and TD						
TD	Hilton et al. 1997 (43)	245 adult travelers	<i>L. rhamnosus</i> GG	1-3 wk	R, DB, PC	Reduced incidence (3.9% vs. 7.4%)
TD	Oksanen et al. 1990 (44)	820 adult travelers	<i>L. rhamnosus</i> GG	1-2 wk	R, DB, PC	Nonsign reduction in incidence
TD	Pozo-Olano et al. 1978 (45)	50 adult travelers	<i>L. acidophilus</i> and <i>L. bulgaricus</i>	8 d	R, DB, PC	No effect on incidence
TD	Black et al. 1989 (46)	92 adult travelers	<i>L. acidophilus</i> , <i>L. bulgaricus</i> , <i>B. bifidum</i> , <i>S. thermophilus</i> mixture	2 wk	R, DB, PC	Reduced incidence (43% vs. 71%)
TD	Katellaris et al. 1995 (47)	282 adult travelers	<i>L. acidophilus</i> or <i>L. fermentum</i>	3 wk	R, DB, PC	No effect on incidence
TD	Kollaritsch et al. 1989 (48)	1231 adult travelers	<i>S. boulardii</i> / <i>L. acidophilus</i>	3 wk	R, DB, PC	Reduced incidence (34% vs. 42%) No effect on incidence
Infectious diarrhea	Pereg et al. 2005 (49)	319 adult travelers	<i>L. casei</i>	8 wk (6 d/wk)	R, SB, PC	No effect on incidence and duration
Repeated infections	Lodinova-Zadnikova 2003 (50)	541 adults (military recruits)	<i>E. coli</i>	colonization at birth	R, C	Reduced incidence after 10 y
7. Probiotics and <i>H. pylori</i> infection						
<i>H. pylori</i> eradication treatment	Canducci et al. 2000 (51)	1714 infants	<i>L. acidophilus</i> LB, dose: 1.5 3 10 ¹⁰	10 d	O, R	Eradication rate ↑
<i>H. pylori</i> eradication treatment	Amuzzi et al. 2001 (52)	60 adults (asymptomatic)	<i>L. rhamnosus</i> GG, dose: 1.2 3 10 ¹⁰	14 d	O, R	Adverse effects ↓
<i>H. pylori</i> eradication treatment	Amuzzi et al. 2001 (53)	120 adults (asymptomatic)	<i>L. rhamnosus</i> GG, dose: 1.2 3 10 ¹⁰	14 d	DB, PC	Adverse effects ↓
<i>H. pylori</i> eradication treatment	Sheu et al. 2002 (54)	160 adults (dyspeptic)	<i>L. acidophilus</i> La5 and <i>B. lactis</i> Bb12, dose: 1 3 10 ¹⁰	4 wk	O, R	Eradication rate ↑ Adverse effects ↓
<i>H. pylori</i> eradication treatment	Cremonini et al. 2002 (55)	85 adults (asymptomatic)	1. <i>L. rhamnosus</i> GG, 2. <i>Saccharomyces boulardii</i> , 3. <i>L. acidophilus</i> La5 and <i>B. lactis</i> Bb12, dose: 1-1.5 3 10 ¹⁰	2 wk	DBPC	Adverse effects ↓ by all probiotic groups
<i>H. pylori</i> eradication treatment	Tursi et al. 2004 (98)	70 adults (dyspeptic with resistant <i>H. pylori</i>)	<i>L. casei</i> ssp. <i>casei</i> DG, dose: 1.6 3 10 ¹⁰	10 d	O, R	Adverse effects ↓
<i>H. pylori</i> eradication treatment	Sykora et al. 2005 (57)	86 children (dyspeptic)	<i>L. casei</i> DN-114 001, dose: 1 3 10 ¹⁰	2 wk	DB, PC	Eradication rate ↑
<i>H. pylori</i> eradication treatment	Myllyluoma et al. 2005 (58)	46 adults (asymptomatic)	Mixture of <i>L. rhamnosus</i> GG, <i>L. rhamnosus</i> LC705, <i>Propionibacterium freudenreichii</i> spp. <i>shermanii</i> JS and <i>B. breve</i> 99), dose 1.3 3 10 ¹¹ first week then, dose: 6.5 3 10 ¹⁰	4 wk	DB, PC	Adverse effects ↓

(Continued)

TABLE 1 Continued

Disease/marker	Reference	Participants	Type of probiotic(s)	Duration	Type of study	Outcome
<i>H. pylori</i> eradication treatment	Lionetti et al. 2006 (59)	40 children (dyspeptic)	<i>L. reuteri</i> ATCC 55730, dose: 1.3 10 ⁸	20 d	DB, PC	Adverse effects ↓
<i>H. pylori</i> eradication treatment	Sheu et al. 2006 (60)	138 adults (asymptomatic with resistant <i>H. pylori</i>)	<i>Lactobacillus</i> La5 and <i>B. lactis</i> Bb12, dose: 4.3 10 ¹⁰	4 wk before treatment	O, R	Urease activity ↓ at pretreatment, Eradication rate ↑ Adverse effects ↓
<i>H. pylori</i> eradication treatment	Cindoruk et al. 2007 (61)	124 adults (dyspeptic)	<i>S. boulardii</i>	2 wk	DB, PC	Adverse effects ↓
<i>H. pylori</i> eradication treatment	de Bortoli et al. 2007 (62)	206 adults (asymptomatic)	<i>Lactobacillus</i> spp., <i>Bifidobacterium</i> spp., and <i>Streptococcus thermophilus</i> with lactoferin	7 d	DB, PC	Eradication rate ↑ Adverse effects ↓
Probiotic effects on <i>H. pylori</i> infection	Michetti et al. 1999 (63)	20 adults (asymptomatic)	<i>L. acidophilus</i> (johnsonii) La1	14 d	DB, PC	Urease activity ↓
Probiotic effects on <i>H. pylori</i> infection	Felley et al. 2001 (64)	52 adults (asymptomatic)	<i>L. acidophilus</i> (johnsonii) La1	3 wk	DB, PC	Urease activity ↓, <i>H. pylori</i> colonization ↓, Inflammation and gastritis ↓
Probiotic effects on <i>H. pylori</i> infection	Sakamoto et al. 2001 (65)	31 adults (asymptomatic)	<i>L. gasseri</i> OLL2716, dose: 1.8–2.5 3 10 ⁹	8 wk	PC	Serum pepsinogen ↓, Urease activity ↓ Ineffective
Probiotic effects on <i>H. pylori</i> infection	Wendakoon et al. 2002 (66)	27 adults (asymptomatic)	<i>L. casei</i> 03, <i>L. acidophilus</i> 2412 and <i>L. acidophilus</i> ACD1, dose: 2.8 3 10 ¹¹	30 d	O	Urease activity ↓ by live La1
Probiotic effects on <i>H. pylori</i> infection	Cruchet et al. 2003 (67)	236 children (asymptomatic)	Both living and heat-killed <i>L. acidophilus</i> La1 or <i>L. paracasei</i> ST1, dose: 1 10 ¹⁰	4 wk	DB, PC	<i>H. pylori</i> colonization ↓, Inflammation ↓
Probiotic effects on <i>H. pylori</i> infection	Pantoflickova et al. 2003 (68)	50 adults (asymptomatic)	<i>L. acidophilus</i> (johnsonii) La1, dose: 1.25 3 10 ^{9–10}	16 wk	DB, PC	Urease activity tended to ↓
Probiotic effects on <i>H. pylori</i> infection	Cats et al. 2003 (69)	20 adults (asymptomatic), 6 adults in control group	<i>L. casei</i> Shirota, dose: 1.95 3 10 ¹⁰	3 wk	O, C	Urease activity ↓, Gastritis and <i>H. pylori</i> colonization ↓
Probiotic effects on <i>H. pylori</i> infection	Wang et al. 2004 (70)	70 dyspeptic adults, endoscopy for 14 participants	<i>L. acidophilus</i> La5 and <i>B. lactis</i> Bb12, dose: 1.3 10 ¹⁰	4 wk	O, C	Eradication ↑ <i>S. boulardii</i> with inulin more effective than <i>L. acidophilus</i> LB
Probiotic effects on <i>H. pylori</i> infection	Gotteland et al. 2005 (71)	254 children (asymptomatic)	<i>L. acidophilus</i> LB or <i>Saccharomyces boulardii</i> (Sb) with inulin, dose: LB 1 310 ¹⁰ , Sb 500 mg + 10 g inulin	8 wk	O, R	Serum gastrin-17 ↓ Urease activity ↓
Probiotic effects on <i>H. pylori</i> infection	Mylyluoma et al. 2007 (101)	16 adults referred for endoscopy	Mixture of <i>L. rhamnosus</i> GG, <i>L. rhamnosus</i> LC705, <i>Propionibacterium freudenreichii</i> spp. <i>shermanii</i> JS and <i>B. lactis</i> BB12, dose: 2.3 10 ¹⁰	8 wk	O, C	Eradication ↑ with both La1 and granberry juice
Probiotic effects on <i>H. pylori</i> infection	Gotteland et al. 2008 (73)	295 children (asymptomatic)	Both living and heat killed <i>L. acidophilus</i> La1 with and without cranberry juice, dose: 8.3 10 ⁸	3 wk	DB, PC	Urease activity ↓ Faecal <i>H. pylori</i> antigen ↓ GSRS improved
Probiotic effects on <i>H. pylori</i> infection	FrancaVilla et al. 2008 (74)	40 adults (dyspeptic)	<i>L. reuteri</i> ATCC 55730	4 wk	DB, PC	

¹ Ab, during antibiotic treatment; CI, confidence interval; CONT, control group; DB, double-blind; EXP, experimental group; NA, not available; O, open; PC, placebo-controlled; R, randomized; RCT, randomized control trial; WMD, weighted mean difference (negative values indicate that duration of diarrhea was shorter in the probiotic than control group); ↑, increase; ↓, decrease.

efficient in the reduction of risk and treatment of diarrhea associated with *C. difficile* infection remains unclear, particularly in the pediatric population.

Additional well-conducted clinical studies using validated clinical outcomes are recommended to further identify populations at high risk of AAD that would benefit most from probiotic intake and to select relevant probiotic candidates. Such validated outcomes should include a conservative definition of diarrhea to differentiate between clinically important cases and negligible changes in stool frequency and/or consistency. In addition, trials should evaluate potential important consequences of AAD such as the need for discontinuation of antibiotic treatment, hospitalization, or intravenous rehydration.

Traveler's diarrhea. TD is a common illness among travelers to developing countries and affects >50% of travelers. Major symptoms involve diarrhea, abdominal cramps, and nausea, and in most cases, bacterial pathogens are the causative agents. Six articles describe the efficacy of various probiotics (*Saccharomyces boulardii*, various lactobacilli, and 1 mix) in populations at high risk for TD.

Two double-blind placebo-controlled RCTs conclude that capsules or sachets containing *L. rhamnosus* GG (2×10^9 CFU/d for 1–2 wk) might prevent TD. Oksanen et al. (44) showed a near-significant reduction of 11.8% in the incidence of diarrhea in healthy Finnish adult travelers to Turkey compared with placebo, with a differential effect dependent on travel location. Hilton et al. (43) found that the incidence of diarrhea was significantly reduced in healthy adults traveling to developing countries when they were taking *L. rhamnosus* GG compared with a placebo capsule. Other studies applying various lactobacilli (Table 1) in doses ranging from 2×10^9 to 2×10^{11} did not show an effect on TD (47,48,86). Two studies applied *Saccharomyces boulardii* at various dose levels (5×10^9 CFU/d, 1×10^{10} CFU/d, 2×10^{10} CFU/d) and found a significant reduction in the incidence of TD among travelers to hot climates in a dose-dependent manner (48,87). Finally, a mixture of probiotic strains was successfully applied to tourists traveling to Egypt, significantly reducing incidence of TD from 71% in the placebo group to 43% in the treatment group (46).

In a recent meta-analysis, the pooled relative risk indicated that some probiotics can significantly prevent TD (88). However, the heterogeneity of the included studies with respect to probiotic strain, dose and duration of treatment, and travel destination and the unidentified causal agents of TD still hamper a more specific recommendation on the use of probiotics. In addition, very few studies report on the stability of the probiotic products, compliance to treatment, and/or recovery of the strain from fecal samples, and such factors may be ultimately important, especially during travel.

NEC in infants

The rationale for probiotic supplementation of preterms is based on data demonstrating differences in the establishment of the intestinal microbiota in preterm infants. The key features of intestinal microbiota in preterm infants, compared with healthy, full-term infants, are 1) restricted number of species, with typically only 3 bacterial species found at 10 d of age; 2) 3 groups including enterobacteria such as *E. coli* and *Klebsiella* spp., enterococci such as *E. faecalis*, and staphylococci such as *S. epidermidis*, *S. aureus*, and *S. haemolyticus* are the most frequently retrieved; 3) all these facultative anaerobes persist at high levels in the fecal flora of preterm infants; and 4) significantly delayed colonization with anaerobes, especially bifidobacteria

(89–91). Additionally, preterm infants are often cared for at intensive care units and receive a broad-spectrum antibiotic, which further contributes to differences in colonization patterns. The direct consequences of disturbed gut microbiota for overall health are not known, but it has been speculated that abnormal pattern of colonization in preterm infants may contribute to the pathogenesis of NEC and to the increased susceptibility to infections. It has also been suggested that enteral administration of probiotics to preterm newborns could prevent infections and NEC and reduce the use of antibiotics (92).

Two systematic studies aimed at determining the effect of probiotics on the prevention of NEC in preterm infants were performed (Table 1). The first one, reported in 2007, identified 7 RCTs (39). This systematic review found that most of the investigated probiotics might reduce the risk of NEC in preterm neonates with <33 wk gestation. Risk of sepsis did not differ significantly among groups. Similarly, the Cochrane Review (40), published in 2008, found that enteral supplementation of certain probiotics reduced the risk of severe NEC and mortality in preterm infants born <1500 g. There was no evidence of significant reduction of nosocomial sepsis. However, it has been shown in both meta-analyses that not all probiotics tested were equally effective. Most effective were combinations used in the studies by Bin-Nun et al. (93) (*Bifidobacterium infantis* plus *Streptococcus thermophilus* plus *B. bifidus*) and by Lin et al. (94) (*Lactobacillus acidophilus* plus *B. infantis*). One additional RCT documented the efficacy of *L. acidophilus* and *B. bifidum* (42).

The use of probiotics for the prevention of NEC in preterm infants is not a routine practice. Therefore, large, well-designed trials are needed to confirm the results. The efficacy and safety of probiotic supplementation in premature infants, <1000 g, need to be defined.

Helicobacter pylori infection

It has been established that *H. pylori* infection is a major cause of chronic gastritis and peptic ulcer disease and a first-class definite carcinogen for stomach cancer (95). Accordingly, there are still many issues to be solved for reducing *H. pylori* infection and improving outcomes of *H. pylori* infection. Those include improvement of *H. pylori* eradication treatment for prevention of gastritis progression and subsequent gastric cancer development. In the future, the treatment of infection will be made difficult by the rapid rate with which the bacteria acquire resistance to the drugs and also long-term changes in microbiota and undesirable side effects induced by the treatment (96,97). These facts provide the rationale for the use of probiotics in the management of *H. pylori* infection.

As a complement to antibiotics, some probiotics have the potential to improve the eradication rate and the overall tolerance of the treatment (Table 1). The early clinical study by Canducci and coworkers (51) provided evidence that *L. acidophilus* LB improved the eradication rate of *H. pylori* significantly. Since then, numerous studies have shown the ability of various probiotic formulations to decrease side effects of anti-*Helicobacter* treatments such as AAD, epigastric pain, discomfort, and flatulence. *L. rhamnosus* GG was able to reduce the occurrence of eradication treatment side effects in 2 separate studies (52,53). The results for *L. rhamnosus* GG were confirmed later in a study comparing *L. rhamnosus* GG, *Saccharomyces boulardii*, and a combination of *L. acidophilus* and *B. lactis* (55). All groups had beneficial effects during and following an anti-*Helicobacter* treatment. There is also a very recent study confirming the effect for *S. boulardii* (61). Furthermore, the probiotic multispecies combination consisting of *L.*

rhamnosus GG, *L. rhamnosus* LC705, *Propionibacterium freidenreichii* subspecies *shermanii* JS and *B. breve* 99 has also improved tolerance to the standard triple therapy and nonsignificantly increased the eradication rate (58). Also, the combination of *B. animalis* Bb12 and *L. acidophilus* LA5 has been beneficial in 2 different trials (54,60). Furthermore, both *L. casei* ssp. *casei* DG (98) and *L. casei* DN-114 001 (57) supplementation significantly increased the eradication rate, but only the *L. casei* DG was able to alleviate side effects.

There are limited numbers of studies on the ability of probiotics to attenuate changes in gastrointestinal microbiota followed by anti-*Helicobacter* triple treatment. In a pilot study a probiotic combination including 2 strains of *L. acidophilus* (CLT60 and CUL21) and 2 strains of *B. bifidum* (CUL17 and Rhodia) suppressed the rise in the numbers of facultative anaerobes seen in the placebo group (99). Later, the same probiotic product was shown to suppress an increase in antibiotic resistance among enterococci (100). Also, a multispecies probiotic combination (*L. rhamnosus* GG, *L. rhamnosus* LC705, *Propionibacterium freidenreichii* spp. *shermanii* JS, and *B. breve* 99) stabilized microbiota during and following the triple anti-*H. pylori* treatment (72). However, despite the probiotic supplementation, microbiota in all studies were susceptible to the effects of the antibiotics administered to eradicate *H. pylori*.

Probiotics as an alternative to antibiotics have also been the focus of several trials (Table 1). Administration of *L. acidophilus* La1 decreased *H. pylori* density in several different trials (63,64,67,68,73). Also, a decrease in gastric inflammation associated with *H. pylori* infection was evident in 2 of these studies (68,73). However, the regular intake of La1 did not permanently eradicate *H. pylori* in any of the studies. Also, *L. gasseri* OLL2716, *L. casei*, and *L. reuteri* ATCC 55730 were found to be effective in suppression of *H. pylori* and reduction in gastric mucosal inflammation (65,69,74). Similar effects of yogurt containing the combination of *L. acidophilus* La5 and *B. lactis* Bb12 have been reported (70). Recently, a probiotic combination consisting of *L. rhamnosus* GG, *L. rhamnosus* LC705, *Propionibacterium freidenreichii* subspecies *shermanii* JS, and *B. animalis* Bb12 revealed a decrease in gastrin-17 and ¹³C-UBT values with *H. pylori*-infected patients (101). However, not all clinical trials have shown effectiveness. In 1 open study 3 *Lactobacillus* strains were ineffective on *H. pylori* infection (66). However, this study used strains that showed inhibition of *H. pylori* in vitro, but no other probiotic characteristics were documented.

To conclude, there are no consistent beneficial data available on the efficacy of probiotics in *H. pylori* infection. However, there are indeed fewer side effects from anti-*Helicobacter* treatment and the disturbance of the gut microbiota when certain probiotics are used. Also, regular consumption of probiotic products with a specified probiotic strain as an alternative to antibiotics may have some potential in suppression of *H. pylori* infection and gastric inflammation, but more clinical studies are needed to confirm their value.

Respiratory tract infections in adults and children

Thus far only a limited number of studies have addressed the potential effect of probiotics in reducing the risk of common upper respiratory tract infections. No studies were found dealing with management of existing respiratory tract infections. Ten human studies investigating the efficacy of probiotics against airway infections were evaluated, of which 3 were in healthy adults, 3 in athletes/soldiers, 1 in elderly, 1 in infants, and 2 in

children (10–19). A number of these studies investigated the effect of probiotic consumption on both airway and gastrointestinal infections (12,13,17,18).

Each study used a different probiotic strain or mixture (Table 1).

Overall, each of the 10 individual studies reported significant improvements on specific sickness-related outcome parameters, but no overall consistency in the efficacy on particular parameters can be deduced. Some studies show an effect on the overall duration of disease and not on incidence (10,13,19), whereas others report an effect on incidence of the disease but not duration (12,17). Most studies also report effects on symptom scores, although different symptoms were affected in the various studies. The 1 study in infants did not report an improvement on respiratory tract infections.

The variety of probiotic strains that were used complicates the development of full insight into the effect of probiotics on airway infections. Two studies assessing the effect on the duration and severity of the common cold successfully used the same mix of strains (*L. gasseri* PA 16/8, *B. longum* SP 07/3, and *B. bifidum* MF 20/5), but in 1 of the studies they were combined with a vitamin supplement and dosed 10-fold higher (10,11), and some differences in effects on symptoms were noted. Two of the studies investigating probiotic efficacy on airway infections employed *L. rhamnosus* GG (14,17), 1 in marathon runners showing no effect on respiratory symptoms and 1 in children with some effect on respiratory infections. *L. reuteri* ATCC 55730 as well as *B. animalis* Bb-12 have shown no effect on respiratory illness in infants (18), but *L. reuteri* ATCC 55730 was effective in the reduction of absenteeism among workers because of respiratory tract or GI symptoms, although no analysis was performed to differentiate between respiratory tract and GI-related problems (12). Each remaining study employed different probiotic treatments, including *L. casei* DN-114 001, *L. fermentum* VRI 003, and a spore-forming *Bacillus clausii* strain, with varying degrees of success.

The rationale for the choice of the particular probiotics has not been described in all studies. Immunomodulation, efficacy in animal studies, and efficacy in human studies with respect to other conditions such as GI infections seem to underlie choices for *L. rhamnosus* GG and *L. reuteri* ATCC 55730. The use of *B. clausii* spores was based on the observed Th-1-stimulating capacity of *B. clausii* in allergic children. However, this immunomodulating potential was not assessed during the study showing positive effect on the duration of recurrent respiratory tract infections in otherwise healthy children (19), making it impossible to directly correlate the positive clinical outcome to an underlying mechanism. The choice of the mix of *L. gasseri* PA 16/8, *B. longum* SP 07/3, and *B. bifidum* MF 20/5 that were used in 2 common cold studies (10,11) was based on their efficacy to protect mice from intestinal infections with pathogenic *E. coli*, but the relevance of this characteristic to their application in common cold studies is unclear.

Most studies that investigate the effect of probiotics on airway infections hypothesize that stimulation of the immune response may be the underlying mechanism. However, it is still rare that endpoint studies are combined with the investigation of biomarkers that may explain the underlying mechanism. In those studies where biomarkers were assessed alongside primary endpoints, at least some insight may have been obtained in the activity of the particular probiotic strain(s). The studies on common cold prevention with the probiotic mix of *L. gasseri* PA 16/8, *B. longum* SP 07/3, and *B. bifidum* MF 20/5 have shown an increase in the number of both CD4+ and CD8+ lymphocytes

in peripheral blood (10,11), but effects on other immune parameters such as lymphocyte activation and phagocytosis were not found. Furthermore, a study in athletes with *L. fermentum* VRI 003 showed a highly significant reduction in the number of days with respiratory symptoms next to a slight increase in blood-derived IFN- γ (16). These observations may provide some insight in relating immune-modulatory effects, but much more research is needed, especially from studies that integrate the measurements on endpoints with biomarkers of the immune system.

Overall, a promising outlook on the potential of probiotics in the combat of airway infections appears, although different probiotic strains may differ in their effects. The limited number of studies and lack of consistent interventions and populations mean a successful formula cannot yet be distilled.

ENT infections

Very few articles consider the efficacy of oral probiotic intake in ENT infections (Table 1). The effect of a mixture of probiotic strains containing *L. rhamnosus* GG, *L. rhamnosus* LC 705, *B. breve* 99, and *P. freudenreichii* JS on acute otitis media (AOM) was studied by Hatakka et al. (20), who reported that neither the duration nor the incidence of AOM was changed during an intervention period of 6 mo.

In contrast, an unexpected increase in the presence of the pathogen *M. catarrhalis* was found in the probiotic group. In another study by Hatakka et al. (17), the number of children with complications of airway infections (AOM, sinusitis, bronchitis, and pneumonia) was significantly reduced on intake of *L. rhamnosus* GG, although the size of the effect decreased after age adjustment.

An interesting alternative approach to the classic oral intake of probiotics for management of ENT infections is the use of airway commensals applied through nasal or oral sprays. A few multicenter studies showed that a 10-d course of spraying with Group A streptococci applied after standard antibiotic therapy significantly reduces the recurrence rate of pharyngo-tonsillitis in adults and children (22–24), as well as otitis media in children (21). This approach seems very promising, but all the studies were performed by the same research group, so more independent studies are needed.

The mechanisms underlying these types of probiotic interventions are thought to relate to colonization resistance. Studies in individuals prone to otitis, sinusitis, and tonsillitis have shown a microbial disbalance resulting in relatively more potential pathogens and fewer protective bacteria with interfering capacity such as Group A streptococci (102). In addition, antibiotic treatment often reinforces this disbalance. Bacterial interference with commensal Group A streptococci may restore balance in the airway microbiota, ensuring competition with potential pathogens. A factor contributing to success in the recolonization with commensal Group A streptococci may have been the preceding antibiotic treatment in patients suffering from ENT infections.

Infectious complications in surgical and critically ill patients

A category of patients who could potentially benefit most from the health-promoting potential of probiotics in terms of prevention of infections are critically ill patients, especially those admitted to the intensive care unit. In this category of patients the infection risk is very high, as is the associated mortality. A number of studies have been performed with critically ill patients in which positive results, but also negative

results, have been reported. As with other applications, many factors may influence the outcome of intervention in this patient category including use of multi- or monospecies probiotics, amount and type of probiotics administered, methods of administration, the time point when instituted, administration of fibers, and type of tube feeding (103).

Several studies have investigated the potential beneficial effects of probiotics in patients undergoing major abdominal surgery. The first one is from the group of Rayes (104). Patients were divided into 3 groups: a conventional group, a placebo group, and a group that received *Lactobacillus plantarum* 299v. Administration of live lactobacilli did not improve the infection rate in the entire study population but did show a trend toward fewer infections in patients with gastric and pancreas resections. The second study from the same study group, also performed in 2002 with *L. plantarum* 299v, involved patients scheduled for liver transplantation; patients were again divided into 3 groups. In liver transplantation patients this probiotic reduced the incidence of infections, the amount of antibiotics administered, and the length of hospital stay. A mixture of different probiotics (*Pediococcus pentosaceus*, *Leuconostoc mesenteroides*, *Lactobacillus paracasei* ssp. *paracasei* F19, and *L. plantarum* 2362) combined with 4 bioactive fibers was also effective in liver transplantation patients in prevention of postoperative bacterial infections (105). This same mixture of probiotics and bioactive fibers was also used in patients scheduled for pylorus-preserving pancreatoduodenectomy. This study (106) showed that early enteral nutrition with bioactive fibers combined with the mixture of probiotics was able to significantly reduce postoperative bacterial infections in patients following pylorus-preserving pancreatoduodenectomy.

In a study by McNaught et al. (107), patients undergoing major abdominal surgery were randomized to either a control group or a treatment group receiving *L. plantarum* 299v. The administration of the *L. plantarum* 299v did not influence the rate of bacterial translocation, gastric colonization, or incidence of postoperative septic morbidity.

Jain et al. (108) performed a study with patients who had been admitted to an intensive care unit, who received a placebo or a combination of probiotic bacteria (consisting of *L. acidophilus* La5, *Bifidobacterium lactis* Bb12, *Strep. thermophilus*, and *Lactobacillus bulgaricus*) with oligofructose. The administration of this mixture did alter the microbial composition of the upper gastrointestinal tract but had no effect on the intestinal permeability and was not associated with measurable clinical benefits.

Two relatively small and 1 large multicenter trial have addressed the potential for probiotics in preventing infectious complications in acute pancreatitis (AP). AP is an acute inflammatory disease caused by either gallstones or excessive alcohol abuse. AP usually runs a mild, self-limiting course, but ~1 in 5 patients will develop necrotizing pancreatitis, which is associated with a 10–30% mortality rate. Mortality is mostly attributed to infectious complications and infection of (peri) pancreatic necrotic tissue in particular. The infections are thought to be the sequelae of a cascade of events that starts with small-bowel bacterial overgrowth, mucosal barrier failure, and a proinflammatory response leading to translocation of intestinal bacteria. Antibiotic prophylaxis has been shown to be ineffective in preventing these infections. In a first study by Olah et al. (109), fewer patients in the group that received *L. plantarum* 299v developed infections of pancreas necrosis than in the placebo group (1 of 22 vs. 7 of 23). A later study from the same study group, now using a combination of 4 lactobacilli

(*Pediococcus pentosaceus*, *Leuconostoc mesenteroides*, *Lactobacillus paracasei* ssp. *paracasei* F19, and *L. plantarum* 2362) combined with 4 bioactive fibers in a total study group of 62 patients did not reduce the number of infections (110).

The Dutch Acute Pancreatitis Study Group has initiated a multicenter study (PROPATRIA) on the effectiveness of a multispecies mixture of microorganisms on prevention of infectious complications in AP (111). The placebo group was treated with high-fiber, high-energy enteral nutrition administered through a nasogastric tube. The intervention group received the enteral nutrition as well as a daily dose of 10^{10} bacteria: *L. acidophilus*, *Lactobacillus casei*, *Lactobacillus salivarius*, *Lactococcus lactis*, *Bifidobacterium bifidum*, and *Bifidobacterium lactis* for a maximum period of 28 d.

This trial in patients with predicted severe AP, in which a total of 296 patients were included, showed no beneficial effect of the treatment on the occurrence of infectious complications (30% in the intervention group vs. 28% in the placebo group). However, mortality in the intervention group was >2 times higher than that in the placebo group [24 (16%) and 9 (6%), respectively, $P < 0.01$]. In 9 patients in the intervention group (none in the placebo group), bowel ischemia was the cause of death (111).

Animal models: what can be learned and what are the drawbacks?

The immunomodulatory effects of probiotics have been studied extensively in vitro and in animal models. These studies have been of paramount importance in elucidating mechanisms of interaction between probiotics and the immune system. However, for many of the (infectious) diseases for which probiotics are being used or considered, animal models are not available or have little relevance for the human disease.

Animal models are very useful for screening potential candidate probiotic strains before these are used in humans. It is difficult to extrapolate from animal studies to the human target population. This especially holds true for consumer probiotics targeted at risk reduction for infectious diseases in the general population and also for allergic diseases and IBD.

Recommendations and gaps

To maximize learning and to generate data to arrive at more consistent evidence, future investigators designing trials on the effect of probiotics on infectious diseases are recommended to consider the following, in addition to the recommendations as outlined in the accompanying general paper (4).

Include biomarker measurements in clinical endpoint studies. Most human studies addressing the effect of a probiotic intervention on resistance against infectious diseases assess either endpoint parameters, such as episodes of diarrhea or severity of infection-related symptoms, or mechanistic parameters and biomarkers such as microbiota changes or changes in epithelial barrier function or immune function markers. Such designs unfortunately preclude the generation of more insight into underlying mechanisms of probiotic action.

In the broad area of infectious disease, many biomarkers could be measured, and the choice for a particular biomarker depends on several factors. Although general recommendations exist on measurement of biomarkers related to the immune system and intestinal health (112–114), the presumed biological activity of the probiotic (why it was selected) and the physiological mechanisms of the host that are involved in dealing with the infection are additional aspects to take into consideration. For instance, when infectious diarrhea is the focus, it seems more

appropriate to include measures related to colonization resistance such as microbial shifts and fecal pH than when dealing with airway infections. In the latter case, when viruses are involved, it seems appropriate to include CD8+ T lymphocyte and natural-killer cell responses among biomarker measurements. Although general recommendations on which biomarkers to include can not be made, as they depend on the nature of the infection and the biological activity of the probiotic, researchers should take the point into consideration and make efforts to include biomarker measurements in clinical endpoint studies to gain more insight into probiotic mechanisms. This would generate data that may help to assess the usefulness of certain biomarkers in the prediction of endpoint outcomes.

Identification of the infectious agent. As the body's defenses against different types of pathogens are differentiated and specialized, the impact of probiotics on the resistance against various pathogens may depend importantly on the type of probiotic strain used and its specific impact on host defense. The efficacy of a probiotic in the management of a given infection may depend on the 1 hand on its ability to counteract pathogenic mechanisms used by pathogens to invade the host and on the other hand on its capacity to modulate a particular immune function or to compensate for some host risk factors (such as age). The identification of the infectious agent in studies addressing, e.g., airway or gastrointestinal infections may help decipher how different probiotics contribute to the management of infections by the host and why certain probiotic strains may be successful whereas others are not. Ultimately, such knowledge will contribute to the rationale for the selection of a particular probiotic in the combat against a particular infection.

Expected incidence and power calculations. Sample size of the study groups should be based on the primary parameters of the study, and if these are incidence, duration, or severity of an infection, the number of participants should be based on the expected incidence, duration, or severity in that target population. Retrospectively, the incidence or duration of the infection in the (control) study population should be compared with the expected incidence, as this may elucidate study outcomes, especially no-effect findings. In addition, reporting changes in incidences and duration should preferably include absolute differences, and the interpretation of such data should be done in context of what is considered clinically relevant to allow judgment of the impact of the effect.

Confounding factors. Use of medication such as antiviral agents, antibiotics, and immunosuppressants should be carefully reported, as should other possible confounding factors such as breast-feeding, and the stability, compliance, and conditions of use of the probiotic under specific circumstances such as travel. To deal with confounding factors, study treatments should, where possible, be stratified over relevant subgroups. Apart from the above recommendations on design of studies on risk reduction for infections, a number of gaps were identified that could be addressed specifically.

Studies should address the applicability of human models of suboptimal health states [e.g., (physical) stress] that may increase the susceptibility to certain infections, to determine relevance for the general population.

Studies should address the appropriate timing for probiotic intervention in the management of infections, also in relation to long-term probiotic ingestion and possible host adaptation.

Conclusions

The potential of probiotics to contribute to colonization resistance and to modulate immune function parameters offers a clear rationale for the use of probiotics in the management of infectious diseases. The different natures of the infections that have been subject to clinical studies with probiotics obviously prevent any generic conclusions in this area. Furthermore, the lack of consistency among studies focusing on 1 specific infection, in study design, applied probiotic strains, outcome parameters, and study population, along with the still-limited number of studies, often preclude a clear conclusion on the efficacy of probiotics in a specific infectious disease. Moreover, the lack of confirmative studies currently makes it impossible to identify the most promising probiotic strains. Positive exceptions are the management of infectious diarrhea (infants and TD), AAD, and NEC.

To advance our knowledge of the possible working mechanisms of probiotics in infectious diseases and to increase our understanding of the role of various biomarkers in risk reduction of infections, it is essential that more probiotic studies incorporate both clinical outcomes and measurement of biomarkers putatively related to the clinical effect such as immune markers and changes in the microbiota and gut barrier function. Furthermore, study designs should be improved so that the identification of infectious agents when relevant, the choice of the probiotic strain, a retrospective comparison of the actual incidence of infections with the predicted incidence, and a more careful reporting of possible confounders may help to understand the outcome, either positive or negative, of trials assessing the effect of probiotics on infections.

Acknowledgments

D.W., J.M.A., E.M., J.S., H.S., and G.R. wrote, read, and approved the final manuscript. The authors would like to thank past and present scientific project managers from ILSI Europe: Sandra Tuijtelars (also from International Dairy Federation), Carina Madsen, Fiona Samuels, Margarita Corrales, and Agnès Méheust.

Literature Cited

- Nisse S. Ueber die Grundlagen einer neuen ursaechlichen Bekämpfung der pathologischen Darmflora. *Deutsch Med Woch.* 1916;42:1181-4.
- Vollaard EJ, Clasener HA. Colonization resistance. *Antimicrob Agents Chemother.* 1994;38:409-14.
- Wilson M. *Microbial inhabitants of humans—their ecology and role in health and disease.* Cambridge: Cambridge University Press, 2005.
- Rijkers G, Bengmark S, Enck P, Haller D, Herz U, Kalliomaki M, Kudo SE, Lenoir-Wijnkoop I, Mercenier A, et al. Guidance for substantiating the evidence for beneficial effects of probiotics: current status and recommendations for future research. *J Nutr.* 2009;140:671-6.
- Szajewska H, Hoekstra JH, Sandhu B. Management of acute gastroenteritis in Europe and the impact of the new recommendations: a multicenter study. The Working Group on acute Diarrhoea of the European Society for Paediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr.* 2000;30:522-7.
- Szajewska H, Mrukowicz JZ. Probiotics in the treatment and prevention of acute infectious diarrhea in infants and children: a systematic review of published randomized, double-blind, placebo-controlled trials. *J Pediatr Gastroenterol Nutr.* 2001;33: Suppl 2: S17-25.
- Van Niel CW, Feudtner C, Garrison MM, Christakis DA. *Lactobacillus* therapy for acute infectious diarrhea in children: a meta-analysis. *Pediatrics.* 2002;109:678-84.
- Huang JS, Bousvaros A, Lee JW, Diaz A, Davidson EJ. Efficacy of probiotic use in acute diarrhea in children: a meta-analysis. *Dig Dis Sci.* 2002;47:2625-34.
- Allen SJ, Okoko B, Martinez E, Gregorio G, Dans LF. Probiotics for treating infectious diarrhoea. *Cochrane Database Syst Rev.* 2004; CD003048.
- de Vrese M, Winkler P, Rautenberg P, Harder T, Noah C, Laue C, Ott S, Hampe J, Schreiber S, et al. Effect of *Lactobacillus gasseri* PA 16/8, *Bifidobacterium longum* SP 07/3, *B. bifidum* MF 20/5 on common cold episodes: a double blind, randomized, controlled trial. *Clin Nutr.* 2005;24:481-91.
- Winkler P, de Vrese M, Laue C, Schrezenmeier J. Effect of a dietary supplement containing probiotic bacteria plus vitamins and minerals on common cold infections and cellular immune parameters. *Int J Clin Pharmacol Ther.* 2005;43:318-26.
- Tubelius P, Stan V, Zachrisson A. Increasing work-place healthiness with the probiotic *Lactobacillus reuteri*: A randomised, double-blind placebo-controlled study. *Environ Health.* 2005;4:25-29.
- Turchet P, Laurenzano M, Auboiron S, Antoine JM. Effect of fermented milk containing the probiotic *Lactobacillus casei* DN-114001 on winter infections in free-living elderly subjects: a randomised, controlled pilot study. *J Nutr Health Aging.* 2003;7:75-7.
- Kekkonen RA, Vasankari TJ, Vuorimaa T, Haahtela T, Julkunen I, Korpela R. The effect of probiotics on respiratory infections and gastrointestinal symptoms during training in marathon runners. *Int J Sport Nutr Exerc Metab.* 2007;17:352-63.
- Tiollier E, Chennaoui M, Gomez-Merino D, Drogou C, Filaire E, Guezennec CY. Effect of a probiotics supplementation on respiratory infections and immune and hormonal parameters during intense military training. *Mil Med.* 2007;172:1006-11.
- Cox AJ, Pyne DB, Saunders PU, Fricker PA. Oral administration of the probiotic *Lactobacillus fermentum* VRI-003 and mucosal immunity in endurance athletes. *Br J Sports Med.* 2008; doi:10.1136/bjism.2007.044628.
- Hatakka K, Savilahti E, Pönkä A, Meurman JH, Poussa T, Nase L, Saxelin M, Korpela R. Effect of long term consumption of probiotic milk on infections in children attending day care centres: double blind, randomised trial. *BMJ.* 2001;322:1-5.
- Weizman Z, Asli G, Alsheikh A. Effect of a probiotic infant formula on infections in child care centers: comparison of two probiotic agents. *Pediatrics.* 2005;115:5-9.
- Marseglia GL, Tosca M, Cirillo I, Licari A, Leone M, Marseglia A, Castellazzi AM, Ciprandi G. Efficacy of *Bacillus clausii* spores in the prevention of recurrent respiratory infections in children: a pilot study. *Ther Clin Risk Manag.* 2007;3:13-7.
- Hatakka K, Blomgren K, Pohjavuori S, Kaijalainen T, Poussa T, Leinonen M, Korpela R, Pitkaranta A. Treatment of acute otitis media with probiotics in otitis-prone children—a double-blind, placebo-controlled randomised study. *Clin Nutr.* 2007;26:314-21.
- Roos K, Håkansson EG, Holm S. Effect of recolonisation with "interfering" streptococci on recurrences of acute and secretory otitis media in children: randomised placebo controlled trial. *BMJ.* 2001; 322:1-4.
- Roos K, Holm SE, Grahn E, Lind L. Alpha-streptococci as supplementary treatment of recurrent streptococcal tonsillitis: a randomized placebo-controlled study. *Scand J Infect Dis.* 1993;25:31-5.
- Roos K, Holm SE, Grahn-Hakansson E, Lagergren L. Recolonization with selected alpha-streptococci for prophylaxis of recurrent streptococcal pharyngotonsillitis—a randomized placebo-controlled multicentre study. *Scand J Infect Dis.* 1996;28:459-62.
- Falck G, Grahn-Hakansson E, Holm SE, Roos K, Lagergren L. Tolerance and efficacy of interfering alpha-streptococci in recurrence of streptococcal pharyngotonsillitis: a placebo-controlled study. *Acta Otolaryngol.* 1999;119:944-8.
- Szajewska H, Ruszczynski M, Radzikowski A. Probiotics in the prevention of antibiotic-associated diarrhea in children: a meta-analysis of randomized controlled trials. *J Pediatr.* 2006;149:367-72.
- Johnston BC, Supina AL, Vohra S. Probiotics for pediatric antibiotic-associated diarrhea: a meta-analysis of randomized placebo-controlled trials. *CMAJ.* 2006;175:377-83.
- Johnston BC, Supina AL, Ospina M, Vohra S. Probiotics for the prevention of pediatric antibiotic-associated diarrhea. *Cochrane Database Syst Rev.* 2007;CD004827.

28. D'Souza AL, Rajkumar C, Cooke J, Bulpitt CJ. Probiotics in prevention of antibiotic associated diarrhoea: meta-analysis. *BMJ*. 2002;324:1361-6.
29. Cremonini F, Di CS, Nista EC, Bartolozzi F, Capelli G, Gasbarrini G, Gasbarrini A. Meta-analysis: the effect of probiotic administration on antibiotic-associated diarrhoea. *Aliment Pharmacol Ther*. 2002;16:1461-7.
30. Szajewska H, Mrukowicz J. Meta-analysis: non-pathogenic yeast *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhoea. *Aliment Pharmacol Ther*. 2005;22:365-72.
31. Hawrelak JA, Whitten DL, Myers SP. Is *Lactobacillus rhamnosus* GG effective in preventing the onset of antibiotic-associated diarrhoea: a systematic review. *Digestion*. 2005;72:51-6.
32. Sazawal S, Hiremath G, Dhingra U, Malik P, Deb S, Black RE. Efficacy of probiotics in prevention of acute diarrhoea: a meta-analysis of masked, randomised, placebo-controlled trials. *Lancet Infect Dis*. 2006;6:374-82.
33. McFarland LV. Meta-analysis of probiotics for the prevention of antibiotic associated diarrhea and the treatment of *Clostridium difficile* disease. *Am J Gastroenterol*. 2006;101:812-22.
34. Wenus C, Goll R, Loken EB, Biong AS, Halvorsen DS, Florholmen J. Prevention of antibiotic-associated diarrhoea by a fermented probiotic milk drink. *Eur J Clin Nutr*. 2008;62:299-301.
35. Hickson M, D'Souza AL, Muthu N, Rogers TR, Want S, Rajkumar C, Bulpitt CJ. Use of probiotic *Lactobacillus* preparation to prevent diarrhoea associated with antibiotics: randomised double blind placebo controlled trial. *BMJ*. 2007;335:80-4.
36. Ruszczynski M, Radzikowski A, Szajewska H. Clinical trial: effectiveness of *Lactobacillus rhamnosus* (strains E/N, Oxy and Pen) in the prevention of antibiotic-associated diarrhoea in children. *Aliment Pharmacol Ther*. 2008;28:154-61.
37. Szymanski H, Armanska M, Kowalska-Duplaga K, Szajewska H. *Bifidobacterium longum* PL03, *Lactobacillus rhamnosus* KL53A, and *Lactobacillus plantarum* PL02 in the prevention of antibiotic-associated diarrhea in children: a randomized controlled pilot trial. *Digestion*. 2008;78:13-7.
38. Dendukuri N, Costa V, McGregor M, Brophy JM. Probiotic therapy for the prevention and treatment of *Clostridium difficile*-associated diarrhea: a systematic review. *CMAJ*. 2005;173:167-70.
39. Deshpande G, Rao S, Patole S. Probiotics for prevention of necrotizing enterocolitis in preterm neonates with very low birthweight: a systematic review of randomised controlled trials. *Lancet*. 2007;369:1614-20.
40. Alfaleh K, Bassler D. Probiotics for prevention of necrotizing enterocolitis in preterm infants. *Cochrane Database Syst Rev*. 2008; CD005496.
41. Samanta M, Sarkar M, Ghosh P, Ghosh J, Sinha M, Chatterjee S. Prophylactic probiotics for prevention of necrotizing enterocolitis in very low birth weight newborns. *J Trop Pediatr*. 2009;55:128-31.
42. Lin HC, Hsu CH, Chen HL, Chung MY, Hsu JE, Lien RI, Tsao LY, Chen CH, Su BH. Oral probiotics prevent necrotizing enterocolitis in very low birth weight preterm infants: a multicenter, randomized, controlled trial. *Pediatrics*. 2008;122:693-700.
43. Hilton E, Kolakowski P, Singer C, Smith M. Efficacy of *Lactobacillus* GG as a diarrheal preventive in travelers. *J Travel Med*. 1997;4:41-3.
44. Oksanen PJ, Salminen S, Saxelin M, Hamalainen P, Ihantola-Vormisto A, Muurasniemi-Isoviita L, Nikkari S, Oksanen T, Porsti I, Salminen E. Prevention of travellers' diarrhoea by *Lactobacillus* GG. *Ann Med*. 1990;22:53-6.
45. Pozo-Olano J, Warram JHJ, Gomez, RG, Cavazos MG. Effect of a lactobacilli preparation on traveler's diarrhea: A randomized, double blind clinical trial. *Gastroenterology* 1978;74:829-830.
46. Black F, Andersen P, Ørskov J, Ørskov F, Gaarslev K, Laulund S. Prophylactic efficacy of lactobacilli on traveller's diarrhea. *Travel Med*. 1989;7:333-335.
47. Katelaris PH, Salam I, Farthing MJ. Lactobacilli to prevent traveler's diarrhea? *N Engl J Med*. 1995;333:1360-1.
48. Kollaritsch H, Kremsner P, Wiedermann G, Scheiner O. Prevention of traveler's diarrhea: comparison of different non-antibiotic preparations. *Travel Med Int*. 1989;11:9-17.
49. Pereg D, Kimhi O, Tirosh A, Orr N, Kayouf R, Lishner M. The effect of fermented yogurt on the prevention of diarrhea in a healthy adult population. *Am J Infect Control*. 2005;33:122-5.
50. Lodinova-Zadnikova R, Cukrowska B, Tlaskalova-Hogenova H. Oral administration of probiotic *Escherichia coli* after birth reduces frequency of allergies and repeated infections later in life (after 10 and 20 years). *Int Arch Allergy Immunol*. 2003;131:209-11.
51. Canducci F, Armuzzi A, Cremonini F, Cammarota G, Bartolozzi F, Pola P, Gasbarrini G, Gasbarrini A. A lyophilized and inactivated culture of *Lactobacillus acidophilus* increases *Helicobacter pylori* eradication rates. *Aliment Pharmacol Ther*. 2000;14:1625-9.
52. Armuzzi A, Cremonini F, Bartolozzi F, Canducci F, Candelli M, Ojetti V, Cammarota G, Anti M, De LA, et al. The effect of oral administration of *Lactobacillus* GG on antibiotic-associated gastrointestinal side-effects during *Helicobacter pylori* eradication therapy. *Aliment Pharmacol Ther*. 2001;15:163-9.
53. Armuzzi A, Cremonini F, Ojetti V, Bartolozzi F, Canducci F, Candelli M, Santarelli L, Cammarota G, De LA, et al. Effect of *Lactobacillus* GG supplementation on antibiotic-associated gastrointestinal side effects during *Helicobacter pylori* eradication therapy: a pilot study. *Digestion*. 2001;63:1-7.
54. Sheu BS, Wu JJ, Lo CY, Wu HW, Chen JH, Lin YS, Lin MD. Impact of supplement with *Lactobacillus*- and *Bifidobacterium*-containing yogurt on triple therapy for *Helicobacter pylori* eradication. *Aliment Pharmacol Ther*. 2002;16:1669-75.
55. Cremonini F, Di CS, Covino M, Armuzzi A, Gabrielli M, Santarelli L, Nista EC, Cammarota G, Gasbarrini G, Gasbarrini A. Effect of different probiotic preparations on anti-*Helicobacter pylori* therapy-related side effects: a parallel group, triple blind, placebo-controlled study. *Am J Gastroenterol*. 2002;97:2744-9.
56. Tursi A, Brandimarte G, Giorgetti GM, Forti G, Modeo ME, Gigliobianco A. Low-dose balsalazide plus a high-potency probiotic preparation is more effective than balsalazide alone or mesalazine in the treatment of acute mild-to-moderate ulcerative colitis. *Med Sci Monit*. 2004;10:1126-31.
57. Sykora J, Valeckova K, Amlerova J, Siala K, Dedek P, Watkins S, Varvarovska J, Stozicky F, Pazdiora P, Schwarz J. Effects of a specially designed fermented milk product containing probiotic *Lactobacillus casei* DN-114 001 and the eradication of *H. pylori* in children: a prospective randomized double-blind study. *J Clin Gastroenterol*. 2005;39:692-8.
58. Myllyluoma E, Veijola L, Ahlroos T, Tynkkynen S, Kankuri E, Vapaatalo H, Rautelin H, Korpela R. Probiotic supplementation improves tolerance to *Helicobacter pylori* eradication therapy—a placebo-controlled, double-blind randomized pilot study. *Aliment Pharmacol Ther*. 2005;21:1263-72.
59. Lionetti E, Miniello VL, Castellana SP, Magista AM, De CA, Maurogiovanni G, Ierardi E, Cavallo L, Francavilla R. *Lactobacillus reuteri* therapy to reduce side-effects during anti-*Helicobacter pylori* treatment in children: a randomized placebo controlled trial. *Aliment Pharmacol Ther*. 2006;24:1461-8.
60. Sheu BS, Cheng HC, Kao AW, Wang ST, Yang YJ, Yang HB, Wu JJ. Pretreatment with *Lactobacillus*- and *Bifidobacterium*-containing yogurt can improve the efficacy of quadruple therapy in eradicating residual *Helicobacter pylori* infection after failed triple therapy. *Am J Clin Nutr*. 2006;83:864-9.
61. Cindoruk M, Erkan G, Karakan T, Dursun A, Unal S. Efficacy and safety of *Saccharomyces boulardii* in the 14-day triple anti-*Helicobacter pylori* therapy: a prospective randomized placebo-controlled double-blind study. *Helicobacter*. 2007;12:309-16.
62. de Bortoli N, Leonardi G, Ciancia E, Merlo A, Bellini M, Costa F, Mumolo MG, Ricchiuti A, Cristiani F, et al. *Helicobacter pylori* eradication: a randomized prospective study of triple therapy versus triple therapy plus lactoferrin and probiotics. *Am J Gastroenterol*. 2007;102:951-6.
63. Michetti P, Dorta G, Wiesel PH, Brassart D, Verdu E, Herranz M, Felley C, Porta N, Rouvet M, et al. Effect of whey-based culture supernatant of *Lactobacillus acidophilus* (*johnsonii*) La1 on *Helicobacter pylori* infection in humans. *Digestion*. 1999;60:203-9.
64. Felley CP, Cortesey-Theulaz I, Rivero JL, Sipponen P, Kaufmann M, Bauerfeind P, Wiesel PH, Brassart D, Pfeifer A, et al. Favourable effect of an acidified milk (LC-1) on *Helicobacter pylori* gastritis in man. *Eur J Gastroenterol Hepatol*. 2001;13:25-9.
65. Sakamoto I, Igarashi M, Kimura K, Takagi A, Miwa T, Koga Y. Suppressing effect of *Lactobacillus gasseri* OLL 2716 (LG21) on

- Helicobacter pylori* infection in humans. J Antimicrob Chemother. 2001;47:709–10.
66. Wendakoon CN, Thomson AB, Ozimek L. Lack of therapeutic effect of a specially designed yogurt for the eradication of *Helicobacter pylori* infection. Digestion. 2002;65:16–20.
 67. Cruchet S, Obregon MC, Salazar G, Diaz E, Gotteland M. Effect of the ingestion of a dietary product containing *Lactobacillus johnsonii* La1 on *Helicobacter pylori* colonization in children. Nutrition. 2003;19:716–21.
 68. Pantoflickova D, Corthesy-Theulaz I, Dorta G, Stolte M, Isler P, Rochat F, Enslin M, Blum AL. Favourable effect of regular intake of fermented milk containing *Lactobacillus johnsonii* on *Helicobacter pylori* associated gastritis. Aliment Pharmacol Ther. 2003;18:805–13.
 69. Cats A, Kuipers EJ, Bosschaert MA, Pot RG, Vandenbroucke-Grauls CM, Kusters JG. Effect of frequent consumption of a *Lactobacillus casei*-containing milk drink in *Helicobacter pylori*-colonized subjects. Aliment Pharmacol Ther. 2003;17:429–35.
 70. Wang KY, Li SN, Liu CS, Perng DS, Su YC, Wu DC, Jan CM, Lai CH, Wang TN, Wang WM. Effects of ingesting *Lactobacillus*- and *Bifidobacterium*-containing yogurt in subjects with colonized *Helicobacter pylori*. Am J Clin Nutr. 2004;80:737–41.
 71. Gotteland M, Poliak L, Cruchet S, Brunser O. Effect of regular ingestion of *Saccharomyces boulardii* plus inulin or *Lactobacillus acidophilus* LB in children colonized by *Helicobacter pylori*. Acta Paediatr. 2005;94:1747–51.
 72. Myllyluoma E, Ahlroos T, Veijola L, Rautelin H, Tynkkyinen S, Korpela R. Effects of anti-*Helicobacter pylori* treatment and probiotic supplementation on intestinal microbiota. Int J Antimicrob Agents. 2007;29:66–72.
 73. Gotteland M, Andrews M, Toledo M, Munoz L, Caceres P, Anziani A, Wittig E, Speisky H, Salazar G. Modulation of *Helicobacter pylori* colonization with cranberry juice and *Lactobacillus johnsonii* La1 in children. Nutrition. 2008;24:421–6.
 74. Francavilla R, Lionetti E, Castellana SP, Magista AM, Maurogiovanni G, Bucci N, De CA, Indrio F, Cavallo L, et al. Inhibition of *Helicobacter pylori* infection in humans by *Lactobacillus reuteri* ATCC 55730 and effect on eradication therapy: a pilot study. Helicobacter. 2008;13:127–34.
 75. Canani RB, Cirillo P, Terrin G, Cesarano L, Spagnuolo MI, De VA, Albano F, Passariello A, De MG, et al. Probiotics for treatment of acute diarrhoea in children: randomised clinical trial of five different preparations. BMJ. 2007;335:340–5.
 76. Szajewska H, Skorka A, Rusczyński M, Gieruszczak-Bialek D. Meta-analysis: *Lactobacillus* GG for treating acute diarrhoea in children. Aliment Pharmacol Ther. 2007;25:871–81.
 77. Szajewska H, Skorka A, Dylag M. Meta-analysis: *Saccharomyces boulardii* for treating acute diarrhoea in children. Aliment Pharmacol Ther. 2007;25:257–64.
 78. Guarino A, Albano F, Ashkenazi S, Gendrel D, Hoekstra JH, Shamir R, Szajewska H. European Society for Paediatric Gastroenterology, Hepatology, and Nutrition/European Society for Paediatric Infectious Diseases evidence-based guidelines for the management of acute gastroenteritis in children in Europe: executive summary. J Pediatr Gastroenterol Nutr. 2008;46:619–21.
 79. Bartlett JG. Clinical practice. Antibiotic-associated diarrhea. N Engl J Med. 2002;346:334–9.
 80. Turck D, Bernet JP, Marx J, Kempf H, Giard P, Walbaum O, Lacombe A, Rembert E, Toursel F, et al. Incidence and risk factors of oral antibiotic-associated diarrhea in an outpatient pediatric population. J Pediatr Gastroenterol Nutr. 2003;37:22–6.
 81. Elstner CL, Lindsay AN, Book LS, Matsen JM. Lack of relationship of *Clostridium difficile* to antibiotic-associated diarrhea in children. Pediatr Infect Dis. 1983;2:364–6.
 82. Lewis S. Response to the article: McFarland LV. Meta-analysis of probiotics for the prevention of antibiotic-associated diarrhea and the treatment of *Clostridium difficile* disease. Am J Gastroenterol. 2006;101:812–22. Am J Gastroenterol. 2007;102:201–2.
 83. Dendukuri N, Brophy J. Inappropriate use of meta-analysis to estimate efficacy of probiotics. Am J Gastroenterol. 2007;102:201–4.
 84. Buts JP, Corthier G, Delmee M. *Saccharomyces boulardii* for *Clostridium difficile*-associated enteropathies in infants. J Pediatr Gastroenterol Nutr. 1993;16:419–25.
 85. Biller JA, Katz AJ, Flores AF, Buie TM, Gorbach SL. Treatment of recurrent *Clostridium difficile* colitis with *Lactobacillus* GG. J Pediatr Gastroenterol Nutr. 1995;21:224–6.
 86. de Dios Pozo-Olano J, Warram JH Jr, Gomez RG, Cavazos MG. Effect of a lactobacilli preparation on traveler's diarrhea. A randomized, double blind clinical trial. Gastroenterology. 1978;74:829–30.
 87. Kollaritsch H, Holst H, Grobara P, Wiedermann G. [Prevention of traveler's diarrhea with *Saccharomyces boulardii*. Results of a placebo controlled double-blind study.] Fortschr Med. 1993;111:152–6.
 88. McFarland LV. Meta-analysis of probiotics for the prevention of traveler's diarrhea. Travel Med Infect Dis. 2007;5:97–105.
 89. Blakey JL, Lubitz L, Barnes GL, Bishop RF, Campbell NT, Gillam GL. Development of gut colonisation in pre-term neonates. J Med Microbiol. 1982;15:519–29.
 90. Gewolb IH, Schwalbe RS, Taciak VL, Harrison TS, Panigrahi P. Stool microflora in extremely low birthweight infants. Arch Dis Child Fetal Neonatal Ed. 1999;80:F167–73.
 91. Magne F, Suau A, Pochart P, Desjeux JF. Fecal microbial community in preterm infants. J Pediatr Gastroenterol Nutr. 2005;41:386–92.
 92. Caplan MS, Jilling T. Neonatal necrotizing enterocolitis: possible role of probiotic supplementation. J Pediatr Gastroenterol Nutr. 2000;30: Suppl 2:S18–22.
 93. Bin-Nun A, Bromiker R, Wilschanski M, Kaplan M, Rudensky B, Caplan M, Hammerman C. Oral probiotics prevent necrotizing enterocolitis in very low birth weight neonates. J Pediatr. 2005;147: 192–6.
 94. Lin HC, Su BH, Chen AC, Lin TW, Tsai CH, Yeh TF, Oh W. Oral probiotics reduce the incidence and severity of necrotizing enterocolitis in very low birth weight infants. Pediatrics. 2005;115:1–4.
 95. Anonymous. Schistosomes, liver flukes and *Helicobacter pylori*. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Lyon, 7–14 June 1994. IARC Monogr Eval Carcinog Risks Hum. 1994;61:1–241.
 96. Deltenre M, Ntounda R, Jonas C, De KE. Eradication of *Helicobacter pylori*: why does it fail? Ital J Gastroenterol Hepatol. 1998;30: Suppl 3:S326–8.
 97. Perri F, Qasim A, Marras L, O'Morain C. Treatment of *Helicobacter pylori* infection. Helicobacter. 2003;8: Suppl 1:53–60.
 98. Tursi A, Brandimarte G, Giorgetti GM, Modeo ME. Effect of *Lactobacillus casei* supplementation on the effectiveness and tolerability of a new second-line 10-day quadruple therapy after failure of a first attempt to cure *Helicobacter pylori* infection. Med Sci Monit. 2004;10:CR662–6.
 99. Madden JA, Plummer SF, Tang J, Garaiova I, Plummer NT, Herbison M, Hunter JO, Shimada T, Cheng L, Shirakawa T. Effect of probiotics on preventing disruption of the intestinal microflora following antibiotic therapy: a double-blind, placebo-controlled pilot study. Int Immunopharmacol. 2005;5:1091–7.
 100. Plummer SF, Garaiova I, Sarvotham T, Cottrell SL, Le SS, Weaver MA, Tang J, Dee P, Hunter J. Effects of probiotics on the composition of the intestinal microbiota following antibiotic therapy. Int J Antimicrob Agents. 2005;26:69–74.
 101. Myllyluoma E, Kajander K, Mikkola H, Kyronpalo S, Rasmussen M, Kankuri E, Sipponen P, Vapaatalo H, Korpela R. Probiotic intervention decreases serum gastrin-17 in *Helicobacter pylori* infection. Dig Liver Dis. 2007;39:516–23.
 102. Brook I. The role of bacterial interference in otitis, sinusitis and tonsillitis. Otolaryngol Head Neck Surg. 2005;133:139–46.
 103. Bengmark S. Synbiotics to strengthen gut barrier function and reduce morbidity in critically ill patients. Clin Nutr. 2004;23:441–5.
 104. Rayes N, Hansen S, Seehofer D, Muller AR, Serke S, Bengmark S, Neuhaus P. Early enteral supply of fiber and lactobacilli versus conventional nutrition: a controlled trial in patients with major abdominal surgery. Nutrition. 2002;18:609–15.
 105. Rayes N, Seehofer D, Theruvath T, Schiller RA, Langrehr JM, Jonas S, Bengmark S, Neuhaus P. Supply of pre- and probiotics reduces bacterial infection rates after liver transplantation—a randomized, double-blind trial. Am J Transplant. 2005;5:125–30.
 106. Rayes N, Seehofer D, Theruvath T, Mogl M, Langrehr JM, Nussler NC, Bengmark S, Neuhaus P. Effect of enteral nutrition and synbiotics on bacterial infection rates after pylorus-preserving pancreatoduodenectomy: a randomized, double-blind trial. Ann Surg. 2007;246: 36–41.

107. McNaught CE, Woodcock NP, MacFie J, Mitchell CJ. A prospective randomised study of the probiotic *Lactobacillus plantarum* 299V on indices of gut barrier function in elective surgical patients. *Gut*. 2002; 51:827–31.
108. Jain PK, McNaught CE, Anderson AD, MacFie J, Mitchell CJ. Influence of synbiotic containing *Lactobacillus acidophilus* La5, *Bifidobacterium lactis* Bb 12, *Streptococcus thermophilus*, *Lactobacillus bulgaricus* and oligofructose on gut barrier function and sepsis in critically ill patients: a randomised controlled trial. *Clin Nutr*. 2004; 23:467–75.
109. Olah A, Belagyi T, Issekutz A, Gamal ME, Bengmark S. Randomized clinical trial of specific *Lactobacillus* and fibre supplement to early enteral nutrition in patients with acute pancreatitis. *Br J Surg*. 2002; 89:1103–7.
110. Olah A, Belagyi T, Poto L, Romics L Jr, Bengmark S. Synbiotic control of inflammation and infection in severe acute pancreatitis: a prospective, randomized, double blind study. *Hepatogastroenterology*. 2007;54:590–4.
111. Besselink MG, van Santvoort HC, Buskens E, Boermeester MA, van Goor H, Timmerman HM, Nieuwenhuijs VB, Bollen TL, van Ramshorst B, et al. Probiotic prophylaxis in predicted severe acute pancreatitis: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2008;371:651–9.
112. Albers R, Antoine JM, Bourdet-Sicard R, Calder PC, Gleeson M, Lesourd B, Samartin S, Sanderson IR, Van LJ, et al. Markers to measure immunomodulation in human nutrition intervention studies. *Br J Nutr*. 2005;94:452–81.
113. Kussmann M, Blum S. OMICS-derived targets for inflammatory gut disorders: opportunities for the development of nutrition related biomarkers. *Endocr Metab Immune Disord Drug Targets*. 2007;7: 271–87.
114. D’Haens G, Sandborn WJ, Feagan BG, Geboes K, Hanauer SB, Irvine EJ, Lemann M, Marteau P, Rutgeerts P, et al. A review of activity indices and efficacy end points for clinical trials of medical therapy in adults with ulcerative colitis. *Gastroenterology*. 2007;132:763–86.